

SYSTEMIC LUPUS ERYTHEMATOSUS DISEASE ACTIVITY INDEX SELENA MODIFICATION

Physicians Global Assessment _____

0 1 2 3
 None Mild Med Severe

SLEDAI SCORE

Check box: If descriptor is present at the time of visit or in the proceeding 10 days

Wt	Present	Descriptor	Definition
8	<input type="checkbox"/>	Seizure	Recent onset. Exclude metabolic, infectious or drug cause
8	<input type="checkbox"/>	Psychosis	Altered ability to function in normal activity due to severe disturbance in the perception of reality. Include hallucinations, incoherence, marked loose associations, impoverished thought content, marked illogical thinking, bizarre, disorganized, or catatonic behavior. Excluded uremia and drug causes.
8	<input type="checkbox"/>	Organic Brain Syndrome	Altered mental function with impaired orientation, memory or other intelligent function, with rapid onset fluctuating clinical features. Include clouding of consciousness with reduced capacity to focus, and inability to sustain attention to environment, plus at least two of the following: perceptual disturbance, incoherent speech, insomnia or daytime drowsiness, or increased or decreased psychomotor activity. Exclude metabolic, infectious or drug causes.
8	<input type="checkbox"/>	Visual Disturbance	Retinal changes of SLE. Include cytoid bodies, retinal hemorrhages, serious exodate or hemorrhages in the choroids, or optic neuritis. Exclude hypertension, infection, or drug causes.
8	<input type="checkbox"/>	Cranial Nerve Disorder	New onset of sensory or motor neuropathy involving cranial nerves.
8	<input type="checkbox"/>	Lupus Headache	Severe persistent headache: may be migrainous, but must be non-responsive to narcotic analgesia.
8	<input type="checkbox"/>	CVA	New onset of cerebrovascular accident(s). Exclude arteriosclerosis
8	<input type="checkbox"/>	Vasculitis	Ulceration, gangrene, tender finger nodules, periungual, infarction, splinter hemorrhages, or biopsy or angiogram proof of vasculitis
4	<input type="checkbox"/>	Arthritis	More than 2 joints with pain and signs of inflammation (i.e. tenderness, swelling, or effusion).
4	<input type="checkbox"/>	Myositis	Proximal muscle aching/weakness, associated with elevated creatine phosphokinase/adolase or electromyogram changes or a biopsy showing myositis.
4	<input type="checkbox"/>	Urinary Casts	Heme-granular or red blood cell casts
4	<input type="checkbox"/>	Hematuria	>5 red blood cells/high power field. Exclude stone, infection or other cause.
4	<input type="checkbox"/>	Proteinuria	>0.5 gm/24 hours. New onset or recent increase of more than 0.5 gm/24 hours.
4	<input type="checkbox"/>	Pyuria	>5 white blood cells/high power field. Exclude infection.
2	<input type="checkbox"/>	New Rash	New onset or recurrence of inflammatory type rash.
2	<input type="checkbox"/>	Alopecia	New onset or recurrence of abnormal, patchy or diffuse loss of hair.
2	<input type="checkbox"/>	Mucosal Ulcers	New onset or recurrence of oral or nasal ulcerations

2	<input type="checkbox"/>	Pleurisy	Pleuritic chest pain with pleural rub or effusion, or pleural thickening.
2	<input type="checkbox"/>	Pericarditis	Pericardial pain with at least 1 of the following: rub, effusion, or electrocardiogram confirmation.
2	<input type="checkbox"/>	Low Complement	Decrease in CH50, C3, or C4 below the lower limit of normal for testing laboratory.
2	<input type="checkbox"/>	Increased DNA binding	>25% binding by Farr assay or above normal range for testing laboratory.
1	<input type="checkbox"/>	Fever	>38°C. Exclude infectious cause
1	<input type="checkbox"/>	Thrombocytopenia	<100,000 platelets/mm ³
1	<input type="checkbox"/>	Leukopenia	<3,000 White blood cell/mm ³ . Exclude drug causes.

_____ TOTAL SCORE (Sum of weights next to descriptors marked present)

Mild or Moderate Flare <input type="checkbox"/>	Severe Flare <input type="checkbox"/>
<input type="checkbox"/> Change in SLEDAI > 3 points	<input type="checkbox"/> Change in SLEDAI > 12
<input type="checkbox"/> New/worse discoid, photosensitive, profundus, cutaneous vasculitis, bullous lupus <input type="checkbox"/> Nasopharyngeal ulcers <input type="checkbox"/> Pleuritis <input type="checkbox"/> Pericarditis <input type="checkbox"/> Arthritis <input type="checkbox"/> Fever (SLE)	<input type="checkbox"/> New/worse CNS-SLE <input type="checkbox"/> Vasculitis <input type="checkbox"/> Nephritis <input type="checkbox"/> Myositis <input type="checkbox"/> Pk < 60.000 <input type="checkbox"/> Home anemia: Hb <7% or decrease in Hb > 3% Requiring: double prednisone <input type="checkbox"/> Prednisone>0.5 mg/kg/day hospitalization
<input type="checkbox"/> Increase in Prednisone, but not to >0.5 mg/kg/day	<input type="checkbox"/> Prednisone >0.5 mg/kg/day
<input type="checkbox"/> Added NSAID or Plaquenil	<input type="checkbox"/> New Cytoxan, Azathioprine, Methotrexate, Hospitalization (SLE)
<input type="checkbox"/> ≥1.0 Increase in PGA, but not to more than 2.5	<input type="checkbox"/> Increase in PGA to > 2.5